

Entanglement in random pure states: Spectral density and average von Neumann entropy

Santosh Kumar and Akhilesh Pandey

School of Physical Sciences, Jawaharlal Nehru University, New Delhi - 110067, India

E-mail: skumar.physics@gmail.com, ap0700@mail.jnu.ac.in

Abstract. Quantum entanglement plays a crucial role in quantum information, quantum teleportation and quantum computation. The information about the entanglement content between subsystems of the composite system is encoded in the Schmidt eigenvalues. We derive here closed expressions for the spectral density of Schmidt eigenvalues for all three invariant classes of random matrix ensembles. We also obtain exact results for average von Neumann entropy. We find that maximum average entanglement is achieved if the system belongs to the symplectic invariant class.

PACS numbers: 03.65.Ud, 03.67.Mn, 02.10.Yn, 05.45.Mt

1. Introduction

Quantum entanglement serves as the measure of quantum non-local correlations between the subsystems of a composite system. It has attracted a great deal of attention in recent times because of its crucial role in quantum information theory, quantum computation and quantum teleportation [1, 2]. Quantum entanglement has bizarre consequences which have baffled physicists over the years [3, 4, 5]. Despite this, quantum entanglement is now universally acknowledged as a useful resource which can be harnessed for a number of applications, e.g., superdense coding [6, 7], quantum teleportation [8], quantum cryptography [9], creation of a quantum computer [10] etc.

In most cases it is desirable to have maximum possible entanglement between the subsystems of the compound system. It is known that random pure states lead to nearly maximal average entanglement. Most of the work so far has concentrated on random pure states which belong to the unitary invariant class ($\beta = 2$) [11, 12, 13, 14, 15, 16, 17, 18], which corresponds to systems with broken time-reversal symmetry. It is natural to investigate time-reversal symmetric systems, i.e., random pure states belonging to the orthogonal ($\beta = 1$) or symplectic ($\beta = 4$) invariant classes. Recently there has been some work on systems with orthogonal invariance and now there are some important results available for systems with finite as well as large Hilbert space dimensions [19, 20, 21, 22]. For systems with symplectic invariance there has been work concerning only large dimensions and for extreme eigenvalue statistics [19, 23, 24]. For these systems the distribution of Schmidt eigenvalues and various entanglement measures have not been investigated for finite dimensional cases.

Our purpose in this paper is to treat the random pure states belonging to all three invariant classes of random matrices [25, 26] on equal footing and obtain results valid for systems with arbitrary Hilbert space dimensions. We derive here the spectral density of Schmidt eigenvalues for arbitrary dimensions. We also derive the average von Neumann entropy and show that maximum average entanglement is achieved if one considers systems belonging to the symplectic invariant class.

The presentation scheme in this paper is as follows. We begin with the brief introduction to the Wishart-Laguerre ensemble and give exact results for the spectral density in Section 2. In section 3 we briefly discuss the quantum entanglement problem in bipartite systems. Section 4 deals with the fixed-trace Wishart-Laguerre ensemble which is relevant for the entanglement problem. We obtain in this section closed results for the spectral density of Schmidt eigenvalues for all three invariant classes of random matrix ensembles. In section 5 we calculate the average von Neumann entropy. By comparing the von-Neumann entropies in the three cases, we show that, as far as average value is concerned, maximum entanglement is achieved for $\beta = 4$. Finally, we conclude in section 6 with summary and some general remarks. Some proofs of the results are outlined in the appendices.

2. Wishart-Laguerre ensemble

Wishart-Laguerre ensembles were introduced by Wishart [27] in connection with the analysis of multivariate distributions. These ensembles comprise the N -dimensional matrices $\mathbf{H} = \mathbf{X}\mathbf{X}^\dagger$ where the \mathbf{X} are $N \times M$ dimensional matrices having its elements as independent and identical (iid) Gaussian random variables.

Wishart-Laguerre ensembles arise explicitly in a number of problems in completely unrelated areas. Besides the quantum entanglement problem [28, 29, 30] some other examples are chaotic mesoscopic systems [31], low energy QCD and gauge theories [32], quantum gravity [33, 34], quantitative finance [35, 36], communication theory [37, 38], gene expression data analysis [39, 40] etc.

2.1. Joint probability density of eigenvalues

Consider the elements of the matrices \mathbf{X} constituting the Wishart-Laguerre ensemble $\mathbf{X}\mathbf{X}^\dagger$ as real, complex or quaternion-real zero-mean iid Gaussian variables. These three cases lead to the three invariant classes of ensembles referred to as orthogonal, unitary and symplectic ensembles and are designated by the Dyson index $\beta = 1, 2$ and 4 [25, 26]. We choose $N \leq M$ for definiteness. The probability distribution followed by \mathbf{X} is

$$P(\mathbf{X}) \propto \exp\left(-\frac{\mathbf{X}\mathbf{X}^\dagger}{2v^2}\right), \quad (1)$$

where v^2 is the variance of each distinct real component of matrix elements of \mathbf{X} . The respective joint probability density (jpd) of eigenvalues ($x_j \in [0, \infty), j = 1, \dots, N$) of $\mathbf{X}\mathbf{X}^\dagger$ for the three cases come out as [26, 41, 42]

$$P^{(\beta)}(\{x\}) = C_{M,N}^{(\beta)} |\Delta_N(\{x\})|^\beta \exp\left(-\sum_{i=1}^N x_i\right) \prod_{j=1}^N x_j^\omega, \quad (2)$$

where $\{x\}$ denotes $\{x_1, \dots, x_N\}$, $\Delta_N(\{x\}) = \prod_{j < k} (x_j - x_k)$ is the Vandermonde determinant and the parameter ω is defined as

$$\omega = \frac{\beta}{2}(M - N + 1) - 1. \quad (3)$$

The normalization $C_{M,N}^{(\beta)}$ can be evaluated using the Selberg integral [25] and turns out to be

$$C_{M,N}^{(\beta)} = \prod_{j=1}^N \frac{\Gamma(\frac{\beta}{2} + 1)}{\Gamma(\frac{\beta}{2}j + 1)\Gamma(\frac{\beta}{2}(j-1) + \omega + 1)}. \quad (4)$$

Note that (2) is obtained from (1) for the choice $v^2 = 1/2$. Some authors choose $v^2 = 1$ (see for example [43]), in which case apart from the change in normalization, the jpd in (2) will have additional factor of $1/2$ in the exponential.

2.2. Spectral densities for eigenvalues

In many cases one is interested in the study of quantities which are linear statistics on the eigenvalues. These quantities do not contain products of different eigenvalues. The averages of these can be evaluated using a single integral over the level-density of the eigenvalues, defined as

$$R_1^{(\beta)}(x) = N \int_0^\infty \cdots \int_0^\infty P^{(\beta)}(x, x_2, \dots, x_N) dx_2 \dots dx_N. \quad (5)$$

From random matrix theory [25] we know that, for the classical ensembles, the above density can be obtained in terms of classical orthogonal polynomials for $\beta = 2$ and in terms of classical skew-orthogonal polynomials for $\beta = 1, 4$ [44, 45, 46, 47, 48].

For the Wishart-Laguerre ensemble defined by jpd (2), the exact expressions for level-densities are known for all M, N [49] in terms of associated Laguerre polynomials $L_\mu^{(b)}(x)$ [50] and the corresponding weight function $w_b(x) = x^b e^{-x}$, $b > -1$. Consider the parameter a defined by

$$2a + 1 = |M - N|. \quad (6)$$

We have for unitary ensemble ($\beta = 2$),

$$\begin{aligned} R_1^{(2)}(N; a; x) &= w_{2a+1}(x) \sum_{\mu=0}^{N-1} \frac{\Gamma(\mu+1)}{\Gamma(\mu+2a+2)} \left(L_\mu^{(2a+1)}(x) \right)^2 \\ &= w_{2a+1}(x) \frac{\Gamma(N+1)}{\Gamma(N+2a+1)} \left[L_{N-1}^{(2a+1)}(x) L_N^{(2a+2)}(x) - L_N^{(2a+1)}(x) L_{N-1}^{(2a+2)}(x) \right]. \end{aligned} \quad (7)$$

The second line in the above equation has been obtained using the Christoffel–Darboux sum [50]. Other expressions equivalent to the above sum are also possible; see (C.1).

For orthogonal ensemble ($\beta = 1$) the level density as well as higher order correlation functions are given in terms of skew-orthogonal polynomials [44]. These skew-orthogonal polynomials can be expanded in terms of orthogonal polynomials [47, 48]. Closed form expressions for the level density can be obtained using these expansions after some tedious algebra. We obtain for even N [49],

$$\begin{aligned} R_{1,\text{even}}^{(1)}(N; a; x) &= 2R_1^{(2)}(N; a; 2x) \\ &\quad - w_{2a+1}(2x) L_{N-1}^{(2a+1)}(2x) \sum_{\mu=0}^{\frac{N-2}{2}} \frac{\Gamma\left(\frac{N+1}{2}\right) \Gamma(\mu+1)}{2^{2a} \Gamma\left(\frac{N+2a+1}{2}\right) \Gamma(\mu+a+2)} L_{2\mu+1}^{(2a+1)}(2x) \\ &\quad + w_a(x) L_{N-1}^{(2a+1)}(2x) \frac{\Gamma\left(\frac{N+1}{2}\right)}{\Gamma\left(\frac{N+2a+1}{2}\right)} \left[\frac{2\Gamma(a+1, x)}{\Gamma(a+1)} - 1 \right], \end{aligned} \quad (8)$$

while for odd N we get

$$\begin{aligned} R_{1,\text{odd}}^{(1)}(N; a; x) &= 2R_1^{(2)}(N; a; 2x) \\ &\quad - w_{2a+1}(2x) L_{N-1}^{(2a+1)}(2x) \sum_{\mu=0}^{\frac{N-1}{2}} \frac{\Gamma\left(\frac{N+1}{2}\right) \Gamma(\mu+\frac{1}{2})}{2^{2a} \Gamma\left(\frac{N+2a+1}{2}\right) \Gamma(\mu+a+\frac{3}{2})} L_{2\mu}^{(2a+1)}(2x) \\ &\quad + w_a(x) L_{N-1}^{(2a+1)}(2x) \frac{\Gamma\left(\frac{N+1}{2}\right)}{\Gamma\left(\frac{N+2a+1}{2}\right)}, \end{aligned} \quad (9)$$

The above two expressions may be put together as a single expression by defining

$$c = N(\bmod 2), \quad (10)$$

so that we have for all N ,

$$\begin{aligned} R_1^{(1)}(N; a; x) &= 2R_1^{(2)}(N; a; 2x) \\ &- w_{2a+1}(2x) L_{N-1}^{(2a+1)}(2x) \sum_{\mu=0}^{\frac{N-2+c}{2}} \frac{\Gamma\left(\frac{N+1}{2}\right) \Gamma\left(\mu + 1 - \frac{c}{2}\right)}{2^{2a} \Gamma\left(\frac{N+2a+1}{2}\right) \Gamma\left(\mu + a + 2 - \frac{c}{2}\right)} L_{2\mu+1-c}^{(2a+1)}(2x) \\ &+ w_a(x) L_{N-1}^{(2a+1)}(2x) \frac{\Gamma\left(\frac{N+1}{2}\right)}{\Gamma\left(\frac{N+2a+1}{2}\right)} \left[(1-c) \frac{2\Gamma(a+1, x)}{\Gamma(a+1)} + (2c-1) \right]. \end{aligned} \quad (11)$$

Here $\Gamma(b, x)$ represents the incomplete Gamma function, $\Gamma(b, x) = \int_x^\infty y^{b-1} e^{-y} dy$.

Finally we consider symplectic ensemble ($\beta = 4$). Similar to $\beta = 1$, the level density in this case also is given in terms of the corresponding skew-orthogonal polynomials. We use the expansion of these skew-orthogonal polynomials given in terms of orthogonal polynomials to obtain the following closed expression for the level-density [49]:

$$\begin{aligned} R_1^{(4)}(N; a; x) &= \frac{1}{2} R_1^{(2)}(2N; 2a + 1/2; x) \\ &- w_{4a+2}(x) L_{2N}^{(4a+2)}(x) \sum_{\mu=0}^{N-1} \frac{\Gamma(N+1) \Gamma(\mu + \frac{1}{2})}{2^{4a+3} \Gamma(N+2a + \frac{3}{2}) \Gamma(\mu + 2a + 2)} L_{2\mu}^{(4a+2)}(x). \end{aligned} \quad (12)$$

Note that in view of definition given by (6), the parameter a assumes only integral or half-integral values (the Wishart case), however, the above results hold for all real $a > -1$ (the Laguerre case).

Comparison between the above theoretical results and corresponding numerical simulations is shown in figure 1. They are in excellent agreement.




Figure 1. Density, $\rho^{(\beta)}(N; a; x) = R_1^{(\beta)}(N; a; x)/N$, of eigenvalues for $\beta = 1, 2, 4$. Solid lines are the theoretical predictions, whereas the symbols represent the simulation results.

For large N , the density is described by the Marčenko-Pastur formula [51, 52],

$$R_1^{(\beta)}(N; a; x) = \begin{cases} \frac{\sqrt{(x_+ - x)(x - x_-)}}{\pi\beta x}, & x_- \leq x \leq x_+, \\ 0, & \text{otherwise.} \end{cases} \quad (13)$$

Here $x_{\pm} = (\beta M/2)(1 \pm \sqrt{N/M})^2$.

3. Bipartite system

Consider a bipartite partition of an NM -dimensional Hilbert space $\mathcal{H}^{(NM)}$ consisting of subsystems \mathbb{A} and \mathbb{B} which span Hilbert spaces $\mathcal{H}_{\mathbb{A}}^{(N)}$ and $\mathcal{H}_{\mathbb{B}}^{(M)}$ such that $\mathcal{H}^{(NM)} = \mathcal{H}_{\mathbb{A}}^{(N)} \otimes \mathcal{H}_{\mathbb{B}}^{(M)}$. For example, \mathbb{A} may be a set of spins and \mathbb{B} can be a heat bath. Without loss of generality we choose $M \geq N$. We take $|\Psi\rangle$ to be a quantum state of the composite system and $|i^{\mathbb{A}}\rangle$, $|\alpha^{\mathbb{B}}\rangle$ as the complete basis of $\mathcal{H}_{\mathbb{A}}^{(N)}$ and $\mathcal{H}_{\mathbb{B}}^{(M)}$ respectively. The state $|\Psi\rangle$ can then be expanded as the linear combination

$$|\Psi\rangle = \sum_{i=1}^N \sum_{\alpha=1}^M g_{i,\alpha} |i^{\mathbb{A}}\rangle \otimes |\alpha^{\mathbb{B}}\rangle. \quad (14)$$

We focus on the situation where $|\Psi\rangle$ is a pure state, so that the density matrix of the composite system is given by $\rho = |\Psi\rangle\langle\Psi|$, with the constraint $\text{Tr}[\rho] = 1$, which is equivalent to $\langle\Psi|\Psi\rangle = 1$.

The density matrix for the pure state $|\Psi\rangle$ can be written as

$$\rho = \sum_{i=1}^N \sum_{\alpha=1}^M \sum_{j=1}^N \sum_{\gamma=1}^M g_{i,\alpha} g_{j,\gamma}^* |i^{\mathbb{A}}\rangle\langle j^{\mathbb{A}}| \otimes |\alpha^{\mathbb{B}}\rangle\langle\gamma^{\mathbb{B}}| \quad (15)$$

The reduced density matrix for subsystem, say \mathbb{A} , can be obtained by tracing out the subsystem \mathbb{B} . We obtain

$$\rho_{\mathbb{A}} = \sum_{i=1}^N \sum_{j=1}^N W_{i,j} |i^{\mathbb{A}}\rangle\langle j^{\mathbb{A}}|, \quad (16)$$

where $W_{i,j} = \sum_{\alpha=1}^M g_{i,\alpha} g_{j,\alpha}^*$ can be viewed as the matrix elements of some matrix $\mathbf{W} = \mathbf{G}\mathbf{G}^\dagger$, \mathbf{G} being $N \times M$ dimensional. In the diagonal basis, (16) can be written as

$$\rho_{\mathbb{A}} = \sum_{i=1}^N \lambda_i |\lambda_i^{\mathbb{A}}\rangle\langle\lambda_i^{\mathbb{A}}|. \quad (17)$$

The λ_i are referred to as Schmidt eigenvalues. The trace condition restricts the Schmidt eigenvalues to the following constraint

$$\sum_{i=1}^N \lambda_i = \text{tr}(\mathbf{G}\mathbf{G}^\dagger) = 1. \quad (18)$$

$|\Psi\rangle$ is called a random pure state if the coefficients $g_{i,\alpha}$ are chosen at random from some given probability distribution.

4. Fixed trace Wishart-Laguerre ensembles

As discussed in the previous section, the Schmidt eigenvalues correspond to Wishart type of matrices. However, the conservation of probabilities for quantum states constrains the trace of these matrices. In earlier works Wishart-Laguerre ensembles have been used to model random quantum states [29, 30]. However, they do not provide a complete quantitative description of the random quantum states due to the neglect of fixed trace condition [30]. The deviations are prominent particularly when the Hilbert space-dimension of the smaller subsystem is not large. Detailed discussion of superiority of fixed-trace ensemble over the ordinary Wishart-Laguerre ensemble for describing random quantum states can be found in [17, 18].

4.1. Joint probability density of Schmidt eigenvalues

Consider the matrix elements $g_{i,\alpha}$ as iid (real, complex or quaternion-real) Gaussian variables with zero mean, such that the fixed trace condition (18) is satisfied. The probability distribution for \mathbf{G} is given by

$$\mathcal{P}(\mathbf{G}) \propto \delta(\text{tr}(\mathbf{G}\mathbf{G}^\dagger) - 1). \quad (19)$$

Correspondingly the jpd of Schmidt eigenvalues ($\lambda_j \in [0, 1], j = 1, \dots, N$) is obtained as

$$\mathcal{P}^{(\beta)}(\{\lambda\}) = \mathcal{C}_{M,N}^{(\beta)} \delta\left(\sum_{i=1}^N \lambda_i - 1\right) |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega. \quad (20)$$

Note that, with the fixed trace condition, (18), the exponential term in (2) becomes a constant and one is left with (20). The normalization constant in the above equation is

$$\mathcal{C}_{M,N}^{(\beta)} = \Gamma\left(\frac{\beta MN}{2}\right) \mathcal{C}_{M,N}^{(\beta)}, \quad (21)$$

where $\mathcal{C}_{M,N}^{(\beta)}$ is given by (4); see Appendix A. The three β values (1, 2, 4) in (20) again correspond to the cases when the coefficients $g_{i,\alpha}$ in (15) are real, complex or quaternion-real. Also, the choice of variance of Gaussian elements is immaterial in this case.

4.2. Spectral densities for Schmidt eigenvalues

The level-density of Schmidt eigenvalues is,

$$\mathcal{R}_1^{(\beta)}(M, N; \lambda) = N \int_0^\infty \cdots \int_0^\infty \mathcal{P}^{(\beta)}(\lambda, \lambda_2, \dots, \lambda_N) d\lambda_2 \dots d\lambda_N. \quad (22)$$

Note that because of the presence of delta function constraint the original limits of integration $[0, 1]$ can be replaced by $[0, \infty)$.

We give here the exact expressions for the level density of Schmidt eigenvalues obtained from the jpd (20). Proof is briefly outlined in Appendices B and C.

We need the following to express the desired results compactly:

$$A_\mu^{\xi, m, n} = \frac{(-1)^\mu \Gamma(\frac{\xi}{2}m + 1) \Gamma(\frac{\xi}{2}mn)}{\frac{\xi}{2} \Gamma(\mu + 1) \Gamma(\frac{\xi}{2}n - \mu) \Gamma(\frac{\xi}{2}(mn - m + n) - \mu - 1) \Gamma(\mu + \frac{\xi}{2}(m - n) + 2)}, \quad (23)$$

$$B_\mu^{m, n} = \frac{(-1)^\mu 2^{n+1} \Gamma(\frac{mn}{2}) \Gamma(\frac{m+1}{2}) \Gamma(\frac{n+1}{2})}{\pi^{1/2} \Gamma(\frac{m-n+1}{2}) \Gamma(\mu + m - n + 1) \Gamma(\mu + 1) \Gamma(n - \mu)}, \quad (24)$$

$$C_{\mu, \nu}^{m, n} = \frac{2^{m-n-1} \Gamma(\frac{m-n+1}{2}) \Gamma(\nu + \frac{m-n+2-c}{2})}{\Gamma(\frac{mn}{2} - m + n - \mu - 1) \Gamma(\nu + \frac{3-c}{2})} B_\mu^{m, n}, \quad (25)$$

$$D^{m, n} = \frac{2^{m-n} \Gamma(\frac{m+1}{2}) \Gamma(\frac{mn}{2})}{\Gamma(\frac{(m+1)(n-1)}{2}) \Gamma(\frac{n}{2})}, \quad (26)$$

$$K_{\mu, \nu}^{m, n} = \frac{2^{2\mu-2N} (\nu + 2m - 2n + 1) \Gamma(2n - \nu) \Gamma(\mu + \frac{1}{2}) \Gamma(\mu + m - n + \frac{1}{2})}{\Gamma(m + \frac{1}{2}) \Gamma(n + \frac{1}{2}) \Gamma(2\mu - \nu + 1)} A_\nu^{4, m, n}, \quad (27)$$

$$\begin{aligned} \mathcal{S}_\mu(\xi, \eta, b; m, n; y) &= y^{\mu + \frac{\xi}{2}(m-n)} (1 - y)^{-\mu + \frac{\xi}{2}(mn-m+n)-2} \\ &\times {}_2\tilde{F}_1\left(\eta - \frac{\xi}{2}n, \mu - \frac{\xi}{2}(m+1)(n-1); 1 + \frac{\xi}{2}(m-n); \frac{by}{y-1}\right), \end{aligned} \quad (28)$$

$$\mathcal{T}_\mu(\xi; m, n; y) = A_\mu^{\xi, m, n} \left[\frac{\xi}{2} n \mathcal{S}(\mu, \xi, 1, 1; m, n, y) - \left(\frac{\xi}{2} n - \mu - 1 \right) \mathcal{S}(\mu, \xi, 0, 1; m, n; y) \right], \quad (29)$$

$$\begin{aligned} \mathcal{U}_\mu(m, n; y) &= (2y)^{\mu+m-n+1} (1 - 2y)^{-\mu + \frac{mn}{2} - m + n - 1} \\ &\times {}_2\tilde{F}_1\left(1, -\frac{1}{2}(m - n - 1); \frac{mn}{2} - m + n - \mu; 2 - \frac{1}{y}\right). \end{aligned} \quad (30)$$

Here ${}_2\tilde{F}_1(l, m; n; y) = {}_2F_1(l, m; n; y)/\Gamma(c)$ is the regularized hypergeometric function.

We first consider the unitary case ($\beta = 2$), which has been considered by several authors. In [17] the result was announced for the square case ($M = N$) in terms of sum involving hypergeometric ${}_3F_2$. Later the result was extended to rectangular case and was given in terms of sum comprising hypergeometric ${}_5F_4$ [18]. In [21] a triple sum expression has been given. We give here a result, valid for all $M \geq N$, which is much simpler than those obtained earlier and is given as a single sum over the hypergeometric ${}_2F_1$. We have

$$\mathcal{R}_1^{(2)}(M, N; \lambda) = \sum_{\mu=0}^{N-1} \mathcal{T}_\mu(2; M, N; \lambda), \quad (31)$$

The spectral density for the orthogonal case ($\beta = 1$) has been obtained for N even in [21]. We consider here both even and odd N cases. For even N , we obtain

$$\mathcal{R}_{1, \text{even}}^{(1)}(M, N; \lambda) = \left[4 \sum_{\mu=0}^{N-1} \mathcal{T}_\mu\left(4; \frac{M}{2}, \frac{N}{2}; 2\lambda\right) + \sum_{\mu=0}^{N-1} B_\mu^{M, N} \mathcal{U}_\mu(M, N, \lambda) \right]$$

$$\begin{aligned}
& - \sum_{\mu=0}^{N-1} \sum_{\nu=0}^{\frac{N-2}{2}} C_{\mu,\nu}^{M,N} \mathcal{S}_{\mu} \left(4, N - 2\nu - 1, 1; \frac{M}{2}, \frac{N}{2}; 2\lambda \right) \Big] \Theta \left(\frac{1}{2} - \lambda \right) \\
& - D^{M,N} \mathcal{S}_{\frac{-M+N-1}{2}} \left(4, 1, 2; \frac{M}{2}, \frac{N}{2}; \lambda \right),
\end{aligned} \tag{32}$$

while for odd N we have,

$$\begin{aligned}
\mathcal{R}_{1,\text{odd}}^{(1)}(M, N; \lambda) &= \left[4 \sum_{\mu=0}^{N-1} \mathcal{T}_{\mu} \left(4; \frac{M}{2}, \frac{N}{2}; 2\lambda \right) \right. \\
& - \sum_{\mu=0}^{N-1} \sum_{\nu=0}^{\frac{N-1}{2}} C_{\mu,\nu}^{M,N} \mathcal{S}_{\mu} \left(4, N - 2\nu, 1; \frac{M}{2}, \frac{N}{2}; 2\lambda \right) \Big] \Theta \left(\frac{1}{2} - \lambda \right) \\
& + D^{M,N} \mathcal{S}_{\frac{-M+N-1}{2}} \left(4, 1, 2; \frac{M}{2}, \frac{N}{2}; \lambda \right).
\end{aligned} \tag{33}$$

Similar to (11) the two expressions for even and odd cases may be combined into one as,

$$\begin{aligned}
\mathcal{R}_1^{(1)}(M, N; \lambda) &= \left[4 \sum_{\mu=0}^{N-1} \mathcal{T}_{\mu} \left(4; \frac{M}{2}, \frac{N}{2}; 2\lambda \right) + (1 - c) \sum_{\mu=0}^{N-1} B_{\mu}^{M,N} \mathcal{U}_{\mu}(M, N, \lambda) \right. \\
& - \sum_{\mu=0}^{N-1} \sum_{\nu=0}^{\frac{N-2+c}{2}} C_{\mu,\nu}^{M,N} \mathcal{S}_{\mu} \left(4, N - 2\nu - 1 + c, 1; \frac{M}{2}, \frac{N}{2}; 2\lambda \right) \Big] \Theta \left(\frac{1}{2} - \lambda \right) \\
& - (-1)^N D^{M,N} \mathcal{S}_{\frac{-M+N-1}{2}} \left(4, 1, 2; \frac{M}{2}, \frac{N}{2}; \lambda \right),
\end{aligned} \tag{34}$$

$\Theta(y)$ in the above equations is the Heaviside theta function, being 0, 1 respectively for $y < 0$ and $y > 0$.

Finally, we consider the symplectic case ($\beta = 4$). Note that we do not take into account the Kramers degeneracy explicitly and use the jpd given by (20). We obtain

$$\mathcal{R}_1^{(4)}(M, N; \lambda) = \sum_{\mu=0}^{2N-1} \mathcal{T}(4; M, N; \lambda) - \sum_{\mu=0}^{N-1} \sum_{\nu=0}^{2\mu} K_{\mu,\nu}^{M,N} \mathcal{S}_{\nu}(4, 0, 1; M, N; \lambda). \tag{35}$$

Of particular interest is the case of $N = 2$ and arbitrary $M \geq 2$, as it corresponds to a physical situation where a single qubit is entangled to a heat bath. In this case the expression for level density is simple and can be obtained directly from (20). We have for all three β ,

$$\mathcal{R}_1^{(\beta)}(M, 2; \lambda) = \kappa^{(\beta,M)} \lambda^{\frac{\beta}{2}(M-1)-1} (1 - \lambda)^{\frac{\beta}{2}(M-1)-1} |1 - 2\lambda|^{\beta}, \tag{36}$$

where $\kappa^{(\beta,M)}$ is given by

$$\kappa^{(\beta,M)} = \begin{cases} 2^{M-2}(M-1), & \beta = 1, \\ \frac{(M-1)\Gamma(2M)}{(\Gamma(M))^2}, & \beta = 2, \\ \frac{(2M-1)(2M-2)\Gamma(4M)}{6(\Gamma(2M))^2}, & \beta = 4. \end{cases} \tag{37}$$

For large M , λ close to $1/2$ dominates the level density (36) and we obtain,

$$\mathcal{R}_1^{(\beta)}(M, 2; \lambda) \approx \frac{(\beta M)^{\frac{\beta+1}{2}}}{2^{\frac{\beta-3}{2}} \Gamma(\frac{\beta+1}{2})} e^{-\frac{\beta M}{2}(1-2\lambda)^2} |1-2\lambda|^\beta. \quad (38)$$

As can be seen in figure 2 as M increases the two maxima move closer and closer towards $\lambda = 1/2$ and become sharply peaked like delta function for $M \rightarrow \infty$.

We have numerically generated fixed trace Wishart-Laguerre ensemble by considering matrices $\mathbf{X}\mathbf{X}^\dagger/\text{tr}(\mathbf{X}\mathbf{X}^\dagger)$, where $\mathbf{X}\mathbf{X}^\dagger$ as above form the Wishart-Laguerre ensemble. Comparison of the above theoretical results with numerics is shown in figure. They agree perfectly.




Figure 2. Density, $\rho_{\text{FT}}^{(\beta)}(M, N; \lambda) = \mathcal{R}_1^{(\beta)}(M, N; \lambda)/N$, of Schmidt eigenvalues for $\beta = 1, 2, 4$ for several M, N values. Solid lines are the theoretical predictions, whereas the symbols represent the simulation results.

The level densities obtained above give good description of the Schmidt eigenvalues. These expressions are important because the statistical properties of entanglement are encoded in the Schmidt eigenvalues. For the unitary case the one-level density has been found to describe accurately the distribution of the Schmidt eigenvalues over the whole range for coupled quantum systems exhibiting chaos [17]. This confirms the existence of universality for the distribution of the Schmidt eigenvalues in quantum chaos. The level-densities can be used to calculate the averages of quantities which are linear statistics on Schmidt eigenvalues, e.g., purity, von Neumann entropy, Rényi entropy etc., thereby giving insight into the extent of entanglement.

5. Average von Neumann Entropy

As discussed above, the information about the degree of entanglement is stored in the Schmidt eigenvalues. One can construct a suitable measure of entanglement by considering a function of these eigenvalues whose value will tell us how entangled a pure state is. Von Neumann entropy serves as an important measure of degree of entanglement. It is an extension of classical entropy concepts to the field of quantum mechanics. In terms of the Schmidt eigenvalues, it is defined as

$$\mathcal{E}^{(\beta)} = - \sum_{j=1}^N \lambda_j \ln \lambda_j. \quad (39)$$

It acquires the maximum value of $\ln N$, when each of the Schmidt eigenvalues assume the value $1/N$. On the other hand the minimum value of 0 is obtained when one of the eigenvalues is one and rest are zero. These two scenarios correspond respectively to maximally entangled and completely unentangled (separable) cases.

To calculate the average von Neumann entropy, $E^{(\beta)} = \langle \mathcal{E}^{(\beta)} \rangle$, we need to perform the following integral:

$$E^{(\beta)}(M, N) = - \int_0^\infty \cdots \int_0^\infty \left(\sum_{i=1}^N \lambda_i \ln \lambda_i \right) \mathcal{P}^{(\beta)}(\{\lambda\}) d\lambda_1 \dots d\lambda_N. \quad (40)$$

As in (22), we have again considered the limits of integrations as $[0, \infty)$ instead of $[0, 1]$. Von Neumann entropy being a linear statistic, the symmetry of the eigenvalues in jpd allows us to reduce the above average involving N integrals to an average involving a single integral, i.e.,

$$E^{(\beta)}(M, N) = - \int_0^\infty \lambda \ln \lambda \mathcal{R}^{(\beta)}(M, N; \lambda) d\lambda. \quad (41)$$

The above calculation can be done using the results for level density given in preceding section. However, it turns out that the above average over the fixed trace ensemble can be reduced directly to an average over the ordinary Wishart-Laguerre ensemble by introducing an auxiliary gamma function integral [12]. The latter route to calculation of average von-Neumann entropy is somewhat more straightforward and we follow it below.

As shown in Appendix D we have,

$$E^{(\beta)}(M, N) = \psi\left(\frac{\beta MN}{2} + 1\right) - \frac{2}{\beta MN} \int_0^\infty x \ln x R_1^{(\beta)}(N; a; x) dx. \quad (42)$$

where $R_1^{(\beta)}(N; a; x)$ is the level density for ordinary Wishart-Laguerre ensemble, as given in (5) and $\psi(x)$ represents the digamma function defined by

$$\psi(y) = \frac{1}{\Gamma(y)} \int_0^\infty e^{-r} r^{y-1} \ln r dr. \quad (43)$$

Equation (42) gives a compact expression for the average von Neumann entropy for all three β values. For $M \geq N \gg 1$, the average von Neumann entropy can be found from (42) using the Marčenko-Pastur density given by (13). Also in this limit

$$\psi\left(\frac{\beta MN}{2}\right) = \ln\left(\frac{\beta MN}{2}\right) + \mathcal{O}\left(\frac{1}{MN}\right). \quad (44)$$

Using (13), the integral in (42) is obtained as $(\beta MN/2) \ln(\beta M/2) + \beta N^2/4$, thereby giving the large N result for all three β as

$$E^{(\beta)}(M, N) = \ln(N) - \frac{N}{2M}. \quad (45)$$

The above result was obtained for $\beta = 2$ in [12]. See [29] for $\beta = 1$.




Figure 3. Average von Neumann entropy for (a) $N = 2, 3, 4$ and $M = 2$ to 16 (b) $M = 8$ and $N = 2$ to 8.

Exact closed form expressions for the average von Neumann entropy for arbitrary M, N can be calculated using (42) and the results in section 2.2. To evaluate the integral in (42) we use the following identity:

$$\int_0^\infty x^b \ln(x) f(b, x) dx = \frac{\partial}{\partial b} \int_0^\infty x^b f(b, x) dx - \int_0^\infty x^b \frac{\partial f(b, x)}{\partial b} dx. \quad (46)$$

The evaluation of the above integral for the three invariant classes involves the use of properties followed by Laguerre polynomials [50] and its expansion (C.2).

The average von Neumann entropy in $\beta = 2$ case is already known [12, 13]. We have,

$$E^{(2)}(M, N) = \sum_{\mu=M+1}^{MN} \frac{1}{\mu} - \frac{N-1}{2M}. \quad (47)$$

For $\beta = 1$, even N , we obtain

$$E^{(1)} = \psi\left(\frac{MN}{2} + 1\right) - \psi(M) - \frac{(N-1)}{2M} + \frac{(M-1)(N-1)}{MN} \ln 2 + \frac{2^{N-1} \Gamma(\frac{M+1}{2}) \Gamma(\frac{N+1}{2})}{MN \sqrt{\pi}}$$

$$\begin{aligned}
& \times \sum_{\mu=0}^{\frac{N-2}{2}} \sum_{\nu=0}^{N-1} \frac{(-1)^\nu (\nu + M - N + 1) \Gamma(\mu + 1) \psi(\nu + M - N + 1) (-\nu - 1)_{2\mu+1}}{\Gamma(\mu + \frac{M-N+3}{2}) \Gamma(N - \nu) \Gamma(\nu + 1) \Gamma(2\mu + 2)} \\
& - \frac{2^{\frac{M+N+d}{2}} \Gamma(\frac{M+1}{2}) \Gamma(\frac{N+1}{2})}{MN\sqrt{\pi}} \\
& \times \sum_{\mu=1}^{\frac{M-N+d}{2}} \sum_{\nu=0}^{N-1} \frac{(-1)^\nu \Gamma(\mu + \nu + \frac{M-N+2-d}{2}) (\psi(\mu + \nu + \frac{M-N+2-d}{2}) - \ln 2)}{2^\mu \Gamma(\nu + 1) \Gamma(N - \nu) \Gamma(\nu + M - N + 1) \Gamma(\mu + \frac{1-d}{2})} \\
& + \frac{2^M \Gamma(\frac{M+1}{2}) \Gamma(\frac{N+1}{2})}{MN\sqrt{\pi}} \sum_{\nu=0}^{N-1} \frac{(-2)^\nu \Gamma(\nu + \frac{M-N+3}{2}) \psi(\nu + \frac{M-N+3}{2})}{\Gamma(N - \nu) \Gamma(\nu + M - N + 1) \Gamma(\nu + 1)} \\
& + (d - 1) \frac{2^M \Gamma(\frac{M+1}{2}) \Gamma(\frac{N+1}{2})}{MN\sqrt{\pi}} \sum_{\nu=0}^{N-1} \frac{(-2)^\nu \Upsilon(\nu + \frac{M-N+1}{2})}{\Gamma(N - \nu) \Gamma(\nu + M - N + 1) \Gamma(\nu + 1)}. \tag{48}
\end{aligned}$$

Here,

$$(m)_n = \frac{\Gamma(m + n)}{\Gamma(m)} \tag{49}$$

represents the Pochhammer function,

$$d = \text{mod}(M - N, 2), \tag{50}$$

and

$$\Upsilon(k) = \left| \frac{\partial}{\partial \alpha} \left(\frac{\Gamma(2\alpha + 2)}{2^{2\alpha} \Gamma(\alpha + 2)} {}_2F_1(\alpha + 1, \alpha + \frac{3}{2}, \alpha + 2; -1) \right) \right|_{\alpha=k}. \tag{51}$$

For $\beta = 1$, odd N , the result is comparatively simple. We have

$$\begin{aligned}
E^{(1)}(M, N) &= \psi\left(\frac{MN}{2} + 1\right) + \frac{(2M + 2N - MN - 2)}{MN} \psi(M) - \frac{(M + N - 1)}{MN} \psi\left(\frac{M}{2}\right) \\
&+ \frac{(M - 1)(N - 1)}{MN} \ln(2) - \frac{(N^2 - N + 4)}{2MN} \\
&- \frac{2^M \Gamma(\frac{M+1}{2}) \Gamma(\frac{N+1}{2})}{MN\sqrt{\pi}} \sum_{\nu=0}^{N-1} \frac{(-2)^\nu \Gamma(\nu + \frac{M-N+3}{2}) \psi(\nu + M - N + 1)}{\Gamma(N - \nu) \Gamma(\nu + 1) \Gamma(\nu + M - N + 1)}. \tag{52}
\end{aligned}$$

Finally, for $\beta = 4$ we have

$$\begin{aligned}
E^{(4)}(M, N) &= \sum_{\mu=2M+1}^{2MN} \frac{1}{\mu} - \frac{(2N - 1)}{4M} + 2^{2N-2} \Gamma(M) \Gamma(N) \\
&\times \sum_{\mu=0}^{N-1} \sum_{\nu=\mu}^{2N} \frac{(-1)^\nu (\nu + 2M - 2N + 1) (-\nu - 1)_{2\mu} \psi(\nu + 2M - 2N + 2)}{2^{2\mu} \Gamma(\mu + 1) \Gamma(\nu + 1) \Gamma(2N + 1 - \nu) \Gamma(\mu + M - N + 1)}. \tag{53}
\end{aligned}$$

The above expressions can be easily implemented in symbolic manipulation software packages like Mathematica.

For large M, N , (45) derives from (47), (48), (52) and (53) in the following way. In (47) the summation term gives $\ln N$ thereby reproducing (45). In (48) the first four terms along with (44) give back (45). In (52) the first five terms give (45) in the above

M	N	$\beta=1$	$\beta=2$	$\beta=4$
2	2	0.27718	0.33333	0.37063
3	2	0.39772	0.45000	0.48099
3	3	0.59987	0.66230	0.69918
4	2	0.46451	0.50952	0.53490
4	3	0.71324	0.76988	0.80167
4	4	0.86452	0.92240	0.95494
5	2	0.50680	0.54563	0.56692
5	3	0.78503	0.83490	0.86210
5	4	0.96152	1.01441	1.04329
5	5	1.08065	1.13262	1.16108
6	2	0.53595	0.56988	0.58815
6	3	0.83442	0.87844	0.90204
6	4	1.02846	1.07596	1.10143
6	5	1.16366	1.21165	1.23744
6	6	1.26130	1.30789	1.33298

Table 1. Average von Neumann entropy evaluated to 5 decimal places for several $M \geq N$ values.

limit. Similarly in (53) the first two terms lead to (45). For $\beta = 1$ and 4 the remaining summation terms become negligible in this limit.

The average von Neumann entropy for the three invariant cases is shown in figure 3. Table I displays the numerical values for several M, N values. It is clear that maximum average entanglement is obtained for $\beta = 4$.

If for $\beta = 4$ we consider the Kramers degeneracy explicitly, the result for average von Neumann entropy will change to

$$E^{(4)*}(2M, 2N) = E^{(4)}(M, N) + \ln 2. \quad (54)$$

In this case the Hilbert spaces \mathcal{H}_A and \mathcal{H}_B are of dimensions $2N$ and $2M$ respectively with sum of all $2N$ eigenvalues being 1. Again we find $E^{(4)*}(2M, 2N) > E^{(2)}(2M, 2N) > E^{(1)}(2M, 2N)$ and $E^{(4)*}(2M, 2N)/2 > E^{(2)}(M, N) > E^{(1)}(M, N)$.

6. Conclusion

We have given exact results for spectral density and average von Neumann entropy for the random pure states belonging to one of the three invariant classes of random matrix ensembles. We have shown using von Neumann entropy measure that, as far as the average is concerned, maximum entanglement is achieved if one generates states from the symplectic invariant class. Several ways of generating random pure states have been proposed by various authors [29, 30, 53, 54, 55, 56, 57]. It will be of interest to test the theoretical predictions presented here using these and newer methods.

Acknowledgments

The authors would like to thank Arul Lakshminarayan for fruitful discussions. SK acknowledges CSIR India for financial assistance.

Appendix A. Relation between normalization constants in (4) and (21)

First we find out the ratio between the normalizations in the jpd's given by (2) and (20). We have from (20),

$$(\mathcal{C}_{M,N}^{(\beta)})^{-1} = \int_0^\infty \cdots \int_0^\infty \delta\left(\sum_{j=1}^N \lambda_j - 1\right) |\Delta_N(\{\lambda\})|^\beta \prod_{k=1}^N \lambda_k^\alpha. \quad (\text{A.1})$$

We introduce an auxiliary variable t in the above expression and define $G(t)$ as,

$$G(t) = \int_0^\infty \cdots \int_0^\infty \delta\left(\sum_{j=1}^N \lambda_j - t\right) |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega, \quad (\text{A.2})$$

so that $(\mathcal{C}_{M,N}^{(\beta)})^{-1} = G(1)$. Taking the Laplace transform ($t \rightarrow s$) of $G(t)$ we get

$$\tilde{G}(s) = \int_0^\infty \cdots \int_0^\infty |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega e^{-s\lambda_j}. \quad (\text{A.3})$$

We now substitute $s\lambda_j = x_j$ and obtain,

$$\tilde{G}(s) = s^{-\frac{\beta MN}{2}} \int_0^\infty \cdots \int_0^\infty |\Delta_N(\{x\})|^\beta \prod_{j=1}^N x_j^\omega e^{-x_j} = s^{-\frac{\beta MN}{2}} (\mathcal{C}_{M,N}^{(\beta)})^{-1}, \quad (\text{A.4})$$

using (21). Using the inverse Laplace transform result ($s \rightarrow t$)

$$\mathcal{L}^{-1}[s^\alpha] = \frac{t^{-\alpha-1}}{\Gamma(-\alpha)}, \quad (\text{A.5})$$

we obtain

$$G(t) = \frac{t^{MN-1}}{\Gamma(\frac{\beta MN}{2})} (\mathcal{C}_{M,N}^{(\beta)})^{-1}. \quad (\text{A.6})$$

Finally, for $t = 1$ we get

$$\frac{\mathcal{C}_{M,N}^{(\beta)}}{\mathcal{C}_{M,N}^{(\beta)}} = \Gamma\left(\frac{\beta MN}{2}\right). \quad (\text{A.7})$$

Equation (A.7) may also be obtained using the auxiliary gamma function integral method discussed in Appendix D.

Appendix B. Relation between level densities in (5) and (22)

The level density for the Schmidt eigenvalues is (we drop the arguments M, N in the following)

$$\mathcal{R}_1^{(\beta)}(\lambda_1) = N \int_0^\infty d\lambda_2 \cdots \int_0^\infty d\lambda_N \mathcal{C}_N \delta\left(\sum_{j=1}^N \lambda_j - 1\right) |\Delta_N(\{\lambda\})|^\beta \prod_{k=1}^N \lambda_k^\omega. \quad (\text{B.1})$$

We introduce the auxiliary variable t and consider $\mathcal{R}_1^{(\beta)}(\lambda_1; t)$ defined by

$$\mathcal{R}_1^{(\beta)}(\lambda_1; t) = N \int_0^\infty d\lambda_2 \cdots \int_0^\infty d\lambda_N \mathcal{C}_N \delta\left(\sum_{j=1}^N \lambda_j - t\right) |\Delta_N(\{\lambda\})|^\beta \prod_{k=1}^N \lambda_k^\omega. \quad (\text{B.2})$$

Thus, $\mathcal{R}_1^{(\beta)}(\lambda) = \mathcal{R}_1^{(\beta)}(\lambda; 1)$. Laplace transform ($t \rightarrow s$) of the above expression yields

$$\tilde{\mathcal{R}}_1^{(\beta)}(\lambda_1; s) = N \int_0^\infty d\lambda_2 \cdots \int_0^\infty d\lambda_N \mathcal{C}_N |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega e^{-s\lambda_j}. \quad (\text{B.3})$$

This, on using (2) and (5), gives

$$\tilde{\mathcal{R}}_1^{(\beta)}(\lambda; s) = \frac{\mathcal{C}_N}{C_N} \frac{1}{s^{\frac{\beta MN}{2}-1}} R_1^{(\beta)}(s\lambda) = \Gamma\left(\frac{\beta MN}{2}\right) s^{1-\frac{\beta MN}{2}} R_1^{(\beta)}(s\lambda), \quad (\text{B.4})$$

Thus the level density for fixed trace ensemble can be obtained from the level density of ordinary Wishart-Laguerre ensemble by taking inverse Laplace transform ($s \rightarrow t$) of the expression on right hand side of (B.4), and by substituting $t = 1$.

Appendix C. Proofs of equations (31)–(35)

We consider the expression

$$R_1^{(2)}(N; a; x) = w_{2a+1}(x) \frac{\Gamma(N+1)}{\Gamma(N+2a+1)} [L_{N-1}^{(2a+2)}(x) L_{N-1}^{(2a+1)}(x) - L_{N-2}^{(2a+2)}(x) L_N^{(2a+1)}(x)] \quad (\text{C.1})$$

for level density given in (7). Note that $L_j^{(k)}(x) = 0$ for $j < 0$. The above expression (and other equivalent ones) can be obtained from (7) using the results given in [50]. We now use the following standard expansion [50] of Laguerre polynomial in (C.1):

$$L_n^{(b)}(x) = \sum_{\mu=0}^n \frac{\Gamma(n+b+1)(-x)^\mu}{\Gamma(b+\mu+1)\Gamma(n-\mu+1)\Gamma(\mu+1)}. \quad (\text{C.2})$$

This gives an expression for the level density (7) involving double sum. Substituting this expression in (B.4) and performing the inverse Laplace transform using the result

$$\mathcal{L}^{-1}[s^\gamma e^{s\lambda}] = \frac{(t-\lambda)^{-\gamma-1} \Theta(t-\lambda)}{\Gamma(-\gamma)}, \quad (\text{C.3})$$

gives the result for level density of Schmidt eigenvalues. This result is a double sum, however one of the sums can be performed in terms of hypergeometric function defined in (28), thus leaving the final result (31) as a single sum.

To derive (32), (33) we use again (C.1) and the expansion (C.2) for Laguerre polynomial in (8) and (9). Along with (C.3) we also need the following Laplace inverse result:

$$\mathcal{L}^{-1}[s^\gamma e^{s\lambda} \Gamma(\alpha+1, s\lambda)] = (t-2\lambda)^{-\alpha-\gamma-1} \lambda^\alpha {}_2\tilde{F}_1\left(1, -\alpha; -\alpha-\gamma; 2-\frac{t}{\lambda}\right) \Theta(t-2\lambda). \quad (\text{C.4})$$

Equation (35) can similarly be derived from (12).

Appendix D. Auxiliary gamma function intergral method

We introduce an auxiliary gamma function intergral in (40) as

$$\begin{aligned} E^{(\beta)}(M, N) = & -\frac{\mathcal{C}_N}{\Gamma(\Omega)} \int_0^\infty e^{-r} r^{\Omega-1} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N \lambda_k \ln \lambda_k \right) \\ & \times \delta\left(\sum_{i=1}^N \lambda_i - 1\right) |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega d\lambda_1 \dots d\lambda_N dr. \end{aligned} \quad (\text{D.1})$$

This can be reduced to the following after some manipulations:

$$\begin{aligned} E^{(\beta)}(M, N) = & \psi(B) - \frac{\mathcal{C}_N}{\Gamma(\Omega)} \int_0^\infty e^{-r} r^{\Omega-1} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N \lambda_k \ln(r\lambda_k) \right) \delta\left(\sum_{i=1}^N \lambda_i - 1\right) \\ & \times |\Delta_N(\{\lambda\})|^\beta \prod_{j=1}^N \lambda_j^\omega d\lambda_1 \dots d\lambda_N dr. \end{aligned} \quad (\text{D.2})$$

where $\psi(x)$ is the digamma function as defined in (43). Now substituting $\lambda_j = x_j/r$, we obtain

$$\begin{aligned} E^{(\beta)}(M, N) = & \psi(\Omega) - \frac{\mathcal{C}_N}{\Gamma(\Omega)} \int_0^\infty e^{-r} r^{\Omega-1} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N \frac{x_k}{r} \ln x_k \right) \delta\left(\sum_{i=1}^N \frac{x_i}{r} - 1\right) \\ & \times \left| \Delta_N\left(\left\{\frac{x_1}{r}\right\}\right) \right|^\beta \prod_{j=1}^N \left(\frac{x_j}{r}\right)^\omega \frac{dx_1}{r} \dots \frac{dx_N}{r} dr \\ = & \psi(\Omega) - \frac{\mathcal{C}_N}{\Gamma(\Omega)} \int_0^\infty \frac{e^{-r} r^{\Omega-1}}{r^{\frac{\beta MN}{2}}} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N x_k \ln x_k \right) \delta\left(\sum_{i=1}^N x_i - r\right) \\ & \times |\Delta_N(\{x\})|^\beta \prod_{j=1}^N x_j^\omega dx_1 \dots dx_N dr. \end{aligned} \quad (\text{D.3})$$

The choice $\Omega = \beta MN/2 + 1$ gives

$$\begin{aligned}
E^{(\beta)}(M, N) &= \psi\left(\frac{\beta MN}{2} + 1\right) - \frac{\mathcal{C}_N}{\Gamma(\frac{\beta MN}{2} + 1)} \int_0^\infty e^{-r} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N x_k \ln x_k\right) \\
&\quad \times \delta\left(\sum_{i=1}^N x_i - r\right) |\Delta_N(\{x\})|^\beta \prod_{j=1}^N x_j^\omega dx_1 \dots dx_N dr \\
&= \psi\left(\frac{\beta MN}{2} + 1\right) - \frac{\mathcal{C}_N}{\Gamma(\frac{\beta MN}{2} + 1)} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N x_k \ln x_k\right) \\
&\quad \times |\Delta_N(\{x\})|^\beta \prod_{j=1}^N x_j^\omega e^{-x_j} dx_1 \dots dx_N \\
&= \psi\left(\frac{\beta MN}{2} + 1\right) - \frac{1}{\Gamma(\frac{\beta MN}{2} + 1)} \frac{\mathcal{C}_N}{C_N} \int_0^\infty \cdots \int_0^\infty \left(\sum_{k=1}^N x_k \ln x_k\right) P(\{x\}) dx_1 \dots dx_N.
\end{aligned} \tag{D.4}$$

Using the ratio of constants, \mathcal{C}_N/C_N , from (A.7) we eventually obtain (42).

References

- [1] Horodecki R, Horodecki P, Horodecki M and Horodecki K 2009 Quantum entanglement *Rev. Mod. Phys.* **81** 865
- [2] Reid M D, Drummond P D, Bowen W P, Cavalcanti E G, Lam P K, Bachor H A, Andersen U L and Leuchs G 2009 The Einstein-Podolsky-Rosen paradox: From concepts to applications *Rev. Mod. Phys.* **81** 1727
- [3] Einstein A, Podolsky B, Rosen N 1935 Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.* **47** 777
- [4] Schrödinger E 1935 Discussion of probability relations between separated systems *Math. Proc. Camb. Philos. Soc.* **31** 555
- [5] Schrödinger E 1936 Probability relations between separated systems *Math. Proc. Camb. Philos. Soc.* **32** 446
- [6] Bennett C H and Wiesner S J 1992 Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states *Phys. Rev. Lett.* **69** 2881
- [7] Harrow A, Hayden P and Leung D 2004 Superdense coding of quantum states *Phys. Rev. Lett.* **92** 187901
- [8] Bennett C H *et al.* 1993 Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels *Phys. Rev. Lett.* **70** 1895
- [9] Bennet C H and Brassard G 1984, in *Quantum Cryptography: Public Key Distribution and Coin Tossing*, Proceedings of the IEEE International Conference on Computers, Systems & Signal Processing (CS Press), 175
- [10] Nielsen M A and Chuang I L 2000 *Quantum Computation and Quantum Information* (Cambridge University Press).
- [11] Lubkin E 1978 Entropy of an n -system from its correlation with a k -reservoir *J. Math. Phys.* **19** 1028
- [12] Page D N 1993 Average entropy of a subsystem *Phys. Rev. Lett.* **71** 1291
- [13] Sen S 1996 Average entropy of a quantum subsystem *Phys. Rev. Lett.* **77** 1

- [14] Życzkowski K and Sommers H-J 2001 Induced measures in the space of mixed quantum states *J. Phys. A: Math. Gen.* **34** 7111
- [15] Cappellini V, Sommers H-J, and Życzkowski K 2006 Distribution of G concurrence of random pure states *Phys. Rev. A* **74** 062322
- [16] Giraud O 2007 Distribution of bipartite entanglement for random pure states *J. Phys. A: Math. Theor.* **40** 2793
- [17] Kubotani H, Adachi S and Toda M 2008 Exact formula of the distribution of Schmidt eigenvalues for dynamical formation of entanglement in quantum chaos *Phys. Rev. Lett.* **100** 240501
- [18] Adachi S, Toda M and Kubotani H 2009 Random matrix theory of singular values of rectangular complex matrices I: Exact formula of one-body distribution function in fixed-trace ensemble *Ann. Phys.* **324** 2278
- [19] Vivo P, Majumdar S N, and Bohigas O 2007 Large deviations of the maximum eigenvalue in Wishart random matrices *J. Phys. A: Math. Theor.* **40** 4317
- [20] Majumdar S N, Bohigas O, and Lakshminarayan A 2008 Exact minimum eigenvalue distribution of an entangled random pure state *J. Stat. Phys.* **131** 33
- [21] Vivo P 2010 Entangled random pure states with orthogonal symmetry: exact results *J. Phys. A: Math. Theor.* **43** 405206
- [22] Majumdar S N 2010 Extreme eigenvalues of Wishart matrices: Application to entangled bipartite system arXiv:1005.4515
- [23] Nadal C, Majumdar S N, and Vergassola M 2010 Phase transitions in the distribution of bipartite entanglement of a random pure state *Phys. Rev. Lett.* **104** 110501
- [24] Nadal C, Majumdar S N, and Vergassola M 2011 Statistical distribution of quantum entanglement for a random bipartite state *J. Stat. Phys.* **142** 403
- [25] Mehta M L 2004 *Random Matrices* (New York: Academic)
- [26] Forrester P J 2010 *Log-Gases and Random Matrices* (Princeton University Press)
- [27] Wishart J 1928 The generalised product moment distribution in samples from a normal multivariate population *Biometrika* **20A** 32
- [28] Życzkowski K and Sommers H-J 2001 Induced measures in the space of mixed quantum states *J. Phys. A: Math. Gen.* **34** 7111
- [29] Bandyopadhyay J N and Lakshminarayan A 2002 Testing statistical bounds on entanglement using quantum chaos *Phys. Rev. Lett.* **89** 060402
- [30] Kubotani H, Toda M, Adachi S 2006 Universality in dynamical formation of entanglement for quantum chaos *Phys. Rev. A* **74** 032314
- [31] Slevin K and Nagao T 1994 Impurity scattering in mesoscopic quantum wires and the Laguerre ensemble *Phys. Rev. B* **50** 2380
- [32] Verbaarschot J 1994 Spectrum of the QCD Dirac operator and chiral random matrix theory *Phys. Rev. Lett.* **72** 2531
- [33] Ambjorn J, Makeenko Y, and Kristjansen C F 1994 Generalized Penner models to all genera *Phys. Rev. D* **50** 5193
- [34] Akemann G 1997 Universal correlators for multi-arc complex matrix models *Nucl. Phys. B* **507** 475
- [35] Bouchaud J-P and Potters M 2001 *Theory of Financial Risk and Derivative Pricing* (Cambridge: Cambridge University Press).
- [36] Burda Z and Jurkiewicz J 2004 Signal and noise in financial correlation matrices *Physica A* **344** 67
- [37] Telatar I E 1999 Capacity of multi-antenna Gaussian channels *Euro. Trans. Telecommun.* **10** 585
- [38] Kumar S and Pandey A 2010 Random matrix model for Nakagami-Hoyt fading *IEEE Trans. Inf. Th.* **56** 2360
- [39] Alter O, Brown P O, Botstein D 2000 Singular value decomposition for genome-wide expression data processing and modeling *PNAS* **97** 10101
- [40] Holter N S, Mitra M, Maritan A, Cieplak M, Banavar J R and Fedoroff N V 2000 Fundamental patterns underlying gene expression profiles: Simplicity from complexity *PNAS* **97** 8409
- [41] Edelman A 1989 Eigenvalues and condition numbers of random matrices *Ph.D. Dissertation, MIT*

- [42] Kumar S and Pandey A 2009 Universal spectral correlations in orthogonal-unitary and symplectic-unitary crossover ensembles of random matrices *Phys. Rev. E* **79** 026211
- [43] Dumitriu I and Edelman A 2002 Matrix models for beta ensembles *J. Math. Phys.* **43** 5830
- [44] Dyson F J 1972 A class of matrix ensembles *J. Math. Phys.* **13** 90
- [45] Nagao T and Wadati M 1991 Correlation functions of random matrix ensembles related to classical orthogonal polynomials *J. Phys. Soc. Jpn.* **60** 3298
- [46] Adler M, Forrester P J, Nagao T and Moerbeke P van 2000 Classical skew orthogonal polynomials and random matrices *J. Stat. Phys.* **99** 141
- [47] Pandey A and Ghosh S 2001 Skew-orthogonal polynomials and universality of energy-level correlations *Phys. Rev. Lett.* **87** 024102
- [48] Ghosh S and Pandey A 2002 Skew-orthogonal polynomials and random-matrix ensembles *Phys. Rev. E* **65** 046221
- [49] Kumar S and Pandey A 2011 Crossover ensembles of random matrices and skew-orthogonal polynomials *Ann. Phys.*, in press
- [50] Szego G 1975 *Orthogonal polynomials* (Providence, R I: American Mathematical Society).
- [51] Marčenko V A and Pastur L A 1967 Distribution of eigenvalues for some sets of random matrices *Math. USSR-Sbornik* **1** 457
- [52] Vinayak and Pandey A 2010 Correlated Wishart ensembles and chaotic time series *Phys. Rev. E* **81** 036202
- [53] Zanardi P, Zalka C, and Faoro L 2000 Entangling power of quantum evolutions *Phys. Rev. A* **62** 030301(R)
- [54] Scott A J 2004 Multipartite entanglement, quantum-error-correcting codes, and entangling power of quantum evolutions *Phys. Rev. A* **69** 052330
- [55] Giraud O and Georgeot B 2005 Intermediate quantum maps for quantum computation *Phys. Rev. A* **72** 042312
- [56] Weinstein Y S and Hellberg C S 2005 Entanglement generation of nearly random operators *Phys. Rev. Lett.* **95** 030501
- [57] Žnidarič M 2007 Optimal two-qubit gate for generation of random bipartite entanglement *Phys. Rev. A* **76** 012318